

Dietary Antioxidants and Human Diseases

Jitender Sharma^{1*}, Roopali Rajput²

Abstract

Oxidative stress has been implicated in the development of cancer and various chronic inflammatory and degenerative human diseases of cardiovascular, neurological, respiratory and gastrointestinal origin. Dietary antioxidants intake as part of normal diet or supplements can attenuate free radical induced cell damage and reduce the morbidity and mortality associated with these diseases. In this review, role of dietary antioxidants in common oxidative stress associated human diseases will be discussed.

Key words: Oxidative; stress; disease; defense; therapy; protection

1. Introduction

Along with the well-known immune defense system, our body has evolved to harbor an antioxidant defense system that functions to maintain the important redox balance. Oxidative stress that plays a major role in pathogenesis of several clinical conditions involving cardiovascular, respiratory & liver diseases, gastrointestinal & neurological disorders, muscle damage, diabetes, and aging (Pizzino et al 2017), is naturally countered by antioxidants. An antioxidant is defined as a substance that inhibits oxidation. For a better understanding, we can state that any substance that has an oxidizable substrate, and when present in low levels, significantly delays or inhibits the oxidation of that substrate, is an antioxidant (Halliwell and Gutteridge 1995). Owing to essential role of antioxidants, these have deemed interesting candidates to be used in therapy for several diseases (Pizzino et al 2017). In the present review, we aim to summarize the available information on the role of dietary antioxidants in several diseases. This will be relevant to understand the protective role of antioxidants and the implications of our dietary habits.

2. Types of Antioxidants

The types of antioxidants range from those generated endogenously by the body's own cells to exogenous agents such as dietary supplements (Kurutas 2015). Based on the activity, antioxidants can be categorized as enzymatic or non-enzymatic (Flora 2009). The antioxidant enzymes include catalase, superoxide dismutase, and glutathione peroxidase and glutathione reductase and these enzymes aid in repair or elimination of damaged biomolecules. Non-enzymatic antioxidants constitute chain breaking antioxidants, and transition metal binding proteins (Nimse and Pal 2015). Chain breaking antioxidants are again of two types, viz., lipid-phase chain breaking antioxidants and aqueous-phase antioxidants. Vitamin E, carotenoids, flavonoids, and Ubiquinol-10, come under lipid-phase chain breaking antioxidants, whereas aqueous-phase chain breaking antioxidants include Vitamin C, uric acid, albumin-bound bilirubin, protein-bound thiol groups and reduced glutathione. Transition metal binding proteins, like, ferritin, ceruloplasmin, lactoferrin, and transferrin, sequester iron and copper and prevent the production

1. *Corresponding author: Assistant Professor, Department of Biochemistry, All India Institute of Medical Sciences, Bathinda, Punjab-151001, India
E-mail : jitendersharma.clinchem@gmail.com
2. Department of Biochemistry, All India Institute of Medical Sciences, Bathinda, Punjab-151001, India
Formerly at department of Molecular Medicine, National Institute of Tuberculosis Respiratory Diseases, Sr Aurobindo Marg, New Delhi-110030, India

*

of free radicals (Nimse and Pal 2015; Pizzino et al 2017).

3. Sources of dietary antioxidants

Majority of antioxidants are supplied in diet and contain one or more of polyphenols, lipoic and ascorbic acid, carotenoids, resveratrol, lycopene, quercetine, genstein, ellagic acid, ubiquinone and indole-3 carbinole. Several plant-based polyphenolic antioxidants, proanthocyanidins (PACs) from different fruits, vegetables and edible plants, have come into focus for their antioxidant properties (Huang 2018). Apart from these, spices and culinary herbs, such as,

ginger, turmeric and garlic, which are routinely used in our cuisines, have also been regarded as good sources of antioxidants with broad spectrum of health promotion properties (Yashin et al. 2017). For a quick reference, various antioxidants and their sources are presented in the Table 1 and the role of anti-oxidants in different diseased conditions in humans is presented in the Figure 1. It is always good to understand how our dietary habits affect our system. The de-lineation of the molecular mechanisms indirectly re-enforces the age-old maxim, ‘You are what you eat’.




Figure 1: Dietary antioxidants linked with different human diseases.

Table 1. Sources of Natural antioxidants

Antioxidants	Common Sources
Beta-carotene	Carrots, spinach, tomatoes, sweet potatoes, papayas, apricots (Grune et al. 2010)
Vitamin C	Citrus fruits-orange, lemon, bell pepper, broccoli, kale, strawberries, kiwi, pineapple (Chambial et al. 2013)
Vitamin E	Cooking oils-sunflower, safflower, olive, cauliflower, sprouts, broccoli, cereal grains, meat, milk, butter, eggs (Grilo et al. 2014; Rizvi et al. 2014)
Flavonoids polyphenol	Potatoes, tomatoes, lettuce, onions, wheat, dark chocolates, red wine, grapes, black tea (Tsao 2010)
Isoflavones,	Soybean, red clovers, apple (Tsao et al. 2006, Tsao et al. 2003)
Neoflavonoids and Chalcones	
Flavones, Flavonols, Flavanones and Flavanonols	Citrus fruits (Kawaii et al. 1999)
Proanthocyanidins	Grapes, apple, blue berries (Tsao 2010)
Anthocyanidins	Black rice (Anderson et al 2006)
Polyphenolic Amides	Oats, chilli pepper (Davis et al 2007; Bratt et al 2003)
Non-flavonoid	Green tea, blue berries (Tsao 2010)

polyphenols	
Resveratrol	Grapes, red wines (Salehi et al. 2018)
Ellagic acid	Strawberries, raspberries (Abe et al 2012)
Lignan	Flaxseed, sesame (Rodríguez -García et al 2019)
Curcumin	Turmeric (Stanić 2017)
Lycopene	Tomatoes, papaya, watermelon, guava, pink grape-fruit (Emmanouil et al 2016)
Coenzyme Q 10	Organ meat- heart, liver and kidney, fish, wheat bran (Saini 2011)

4. Role of antioxidants in diseases

4.1 Cancer

Several observational studies showed that the risk of cancer is inversely related with dietary vegetables and fruits rich in antioxidants (Willett 2010). In most cancer models, tumor dependent immunosuppression promotes evasion of malignant cells via host's anti-tumor immune responses, which can also be modulated by antioxidants. In a mouse experimental model, antioxidants like curcumin were shown to restore tumor-induced depletion of host CD4+/CD8+ T cell proliferation and inhibition of apoptosis of thymocytes and splenocytes (Bhattacharyya 2010). In cases of breast cancer, Resveratrol, a dietary polyphenol inactivates Stat3, prevents generation and function of tumor-evoked regulatory B cells and inhibits lung metastasis of breast cancer (Lee-Chang et al. 2013). Chemotherapy, radiotherapy and immunotherapy are widely used in the management of majority of cancers. After treatment with these modalities, a sizeable number of patients experience adverse effects due to chemotherapy and/ or radiotherapy induced free radical injury to normal cells. Antioxidants like phytochemical soy isoflavones, protects normal cells from radiotherapy induced toxicity in prostate cancers (Raffoul et al. 2007). Dietary antioxidant supplements like Vitamin C and E reduces risk of breast cancer recurrence and overall mortality from breast cancer (Greenlee 2012).

4.2 Cardiovascular Diseases

After several *in vitro* and *in vivo* studies suggested role of antioxidant vitamins C, E and beta-carotene in preventing or slowing down the progression of atherosclerotic processes (Maeda et al 2000; Devaraj et al 1996; Romanchik et al 1995), large scale clinical trials were conducted. Some experimental studies indicated that antioxidant vitamin

supplementation cause reduction in cardiovascular events by preventing endothelial damage and proliferation and production of foam cells (Stephens et al. 1996; Hozawa et al 2007; Wannamethee et al 2006). Polyphenolic compounds are abundant in plants and are readily found in fruit and vegetables. In addition, they are important components of herbs and spices and are likely to be critical ingredients in Chinese medicines. Polyphenolic compounds can directly interact with ROS, block generation of free radicals by inhibiting xanthine oxidase enzyme and chelates iron and copper (Quideau et al. 2011). Polyphenols prevents lipid peroxidation and uptake of oxidized LDL by macrophage (Yamakoshi et al. 1999; Kaplan et al. 2001). Joshipuræ *et al.* found that intake of green leafy vegetables and a vitamin C rich fruit provides protection against coronary heart diseases (Joshipura et al. 2001) A significant reduction in ischemic stroke, unstable angina, acute myocardial infarction was seen when vitamin E supplements were given to pre-existing coronary heart disease patients (Stephens et al. 1996; Boaz et al. 2000).

4.3 Neurological Disorders

Oxidative stress induced cell damage, impaired damaged DNA repair and mitochondrial dysfunction are well established factors in the development of neurodegenerative disorders like Alzheimer's disease, Parkinson's disease, Huntington's disease and familial amyotrophic lateral sclerosis (Kim et al. 2015; Federico et al. 2012). In addition, increase in protein aggregates activates microglia cells and leads to sustained neuroinflammation (Masgrau et al 2017). Microglia releases several cytokines and chemokines which aggravates oxidative stress and results in neuronal damage (Frank-Cannon et al. 2009). Dietary antioxidants have the potential to prevent, delay, or ameliorate these disorders. In vitro and animal model studies support the potential

beneficial role of various dietary antioxidant compounds in neurodegenerative diseases. In murine models of Alzheimer's disease, dietary lycopene administration attenuates mitochondrial oxidative damage, inhibits pro-inflammatory cytokines in brain and suppress protein aggregation (Prakash and Kumar 2014; Katayama et al 2011).

In Parkinson's disease, lycopene-rich tomato powder intake prevented decline in striatal dopamine neurotransmission and degeneration of nigral dopaminergic neurons in rodent models (Suganuma et al 2002; Di Matteo et al 2009). Similarly, lycopene showed protection against Huntington's disease induced by 3-nitropropionic acid in rodent models (Prakash and Kumar 2009). Also, lycopene treatment was shown to be effective in cultured cell models of Alzheimer's disease and Parkinson's disease (Qu et al. 2011; Yi et al. 2013). Dietary carotenoids like fucoxanthin, astaxanthin, and crocetin were found to be protective against Alzheimer's disease (Xiang et al. 2017; Lobos et al. 2016; Tiribuzi et al. 2017). Astaxanthin inhibits oxidative stress and provides neuroprotection in Parkinson's disease and familial amyotrophic lateral sclerosis (Ye et al. 2013; Isonaka et al. 2011).

4.4 Gastro-intestinal Diseases

Irritable bowel syndrome (IBS) and inflammatory bowel disease (IBD), which includes Crohn's disease and ulcerative colitis are associated with oxidative stress (Balmus et al. 2016; Mete et al. 2013). Several studies suggested role of dietary antioxidants in CD, UC and IBS. Curcumin, derived from *Curcuma longa* plant, found to be useful in ulcerative colitis when combined with mesalamine drug in a random double-blind controlled study (Lang et al. 2015). Other dietary antioxidants, like, boswellic acids (Gerhardt et al 2001), artemesinin, myrcene (Krebs et al 2010), *T. wilfordii* (Zhu et al 2015), and fish oil (Feagan et al 2008), found to be beneficial in Crohn's disease. Also, Chamomile (Langhorst et al 2014), and fish oil (Barbosa et al 2003) improved the oxidative stress in Ulcerative colitis. Liver is the principal detoxifying organ. It metabolizes various compounds that produce reactive oxygen species. Oxidative stress as a result of environmental pollution, drug overdose, alcohol intake and high calorie intake leads to liver diseases (Jadeja et al. 2017). Curcumin and Quercetin has demonstrated hepatoprotective

actions on acute and chronic liver injury (Wang et al 2012; Reyes-Gordillo et al 2007; Hernández-Ortega et al. 2012). Resveratrol protects against alcohol-induced lipid peroxidation in animal models (Kasdallah-Grissa et al. 2006).

4.5 Respiratory Diseases

Vitamin E, vitamin C, beta-carotene, polyphenols, catechins, flavonol and flavone have been speculated to have therapeutic effects in Chronic Obstructive Pulmonary Disease (COPD) patients (Grievink et al. 1998; Santus et al. 2005). Elevated levels of dietary antioxidants lead to lower prevalence of chronic bronchitis and dyspnea (Rautalahti et al. 1997). An increase of Vitamin C levels by 20 mmol / L was associated with a 13% reduction in the risk of COPD (Sargeant et al. 2000). Resveratrol and curcumin inhibit inflammatory response seen in COPD (Kode et al 2008; Biswas et al 2005). Recently, a cross-sectional study in Korean population, revealed that dietary antioxidant like carotene, vitamin A and C intake was beneficial in male smokers with COPD (Hong et al 2018). Oxidative stress might exacerbate asthma by increasing airway inflammation, and responsiveness (Fitzpatrick et al 2009). Several studies suggested an association between low vitamin E intake and increased severity of bronchial asthma (Nurmatov et al 2011; Pearson et al 2004). Vitamin C intake aided recovery in exercise induced asthma (Kurti et al 2016). Carotenoids consumption was found to reduce airway inflammation (Lovett-Racke et al 2002) and associated with low prevalence of asthma in women (Romieu et al. 2006).

5. Conclusion

Oxidative stress occurs from the imbalance between free radical production and antioxidant defenses. Oxidative stress is involved in various diseases such as cancer, cardiovascular, respiratory, gastrointestinal and neurological disorders. Dietary antioxidants are widely reported to exert a protective effect in cells and animal models. Therefore, further research should focus on disease specific, molecular target oriented dietary antioxidants.

ACKNOWLEDGEMENT

JS would like to acknowledge institutional support from AIIMS, Bathinda, Punjab, India.

AUTHOR DECLARATIONS

The authors declare that they have no conflict of interest for this piece of work. JS conceived the

the review article, collected information, prepared the rough draft, proof-read and revised the manuscript, constructed the table and provided final approval for submission of the manuscript. RR provided substantial intellectual input for different sub-sections, created the figure, proof-read and revised the manuscript and approved the final version of the article.

REFERENCES

- Pizzino G, Irrera N, Cucinotta M, et al. (2017). Oxidative Stress: Harms and Benefits for Human Health. *Oxid Med Cell Longev*, 2017:8416763. doi:10.1155/2017/8416763.
- Halliwell B, Gutteridge JC. (1995). The definition and measurement of antioxidants in biological systems. *Free Radic Biol Med*, 18,125–126.
- Kurutas E.B. (2015). The importance of antioxidants which play the role in cellular response against oxidative/nitrosative stress: current state. *Nutr. J*, 71(15),1–22.
- Flora SJ. (2009). Structural, chemical and biological aspects of antioxidants for strategies against metal and metalloid exposure. *Oxid Med Cell Longev*, 2(4), 191–206.
- Nimse SB, Pal D. (2015). Free radicals, natural antioxidants, and their reaction mechanisms. *RSC Adv*, 5, 27986–28006.
- Huang D. (2018). Dietary antioxidants and health promotion. *Antioxidants*, 7(1), 9.
- Yashin A., Yashin Y., Xia X. and Nemzer B. (2017). Antioxidant activity of spices and their impact on human health: a review. *Antioxidants*, 6(3), 70.
- Grune T, Lietz G, Palou A, Ross AC, Stahl W, Tang G, et al. (2010). Beta-carotene is an important vitamin A source for humans *J. Nutr*; 140, 2268S–2285SS.
- Chambial S, Dwivedi S, Shukla KK., John PJ, Sharma P. (2013). Vitamin C in disease prevention and cure: An overview. *Indian J. Clin. Biochem*, 28, 314–328.
- Grilo Câmara E, Costa PN, Gurgel CSS, Beserra AFM, Almeida FNS, Dimenstein R. (2014). Alpha-tocopherol and gamma-tocopherol concentration in vegetable oils. *Food Sci Technol*, 34, 379–385.
- Rizvi S, Raza ST, Ahmed F, Ahmad A, Abbas S, Mahdi F. (2014). The role of vitamin E in human health and some diseases. *Sultan Qaboos Univ. Med. J*, 14, e157–e165.
- Tsao R.. (2010). Chemistry and biochemistry of dietary polyphenols. *Nutrients*, 2, 123– 146.
- Tsao R, Papadopoulos Y, Yang R, Young JC, McRae K. (2006). Isoflavone profiles of red clovers and their distribution in different parts harvested at different growing stages. *J. Agric. Food Chem*, 54, 5797–5805.
- Tsao R, Yang R, Young JC, Zhu H. (2003). Polyphenolic profiles in eight apple cultivars using high-performance liquid chromatography (HPLC). *J. Agric. Food Chem*, 51, 6347–6353.
- Kawaii S, Tomono Y, Katase E, Ogawa K, Yano M. (1999). Quantitation of flavonoid constituents in citrus fruits. *J. Agric. Food Chem*, 47, 3565–3571.
- Anderson OM, Jordheim M. (2006) The anthocyanins.. In: Anderson O.M., Markham K.R., (Ed.), *Flavonoids: Chemistry, Biochemistry and Applications*(pp. 472–551). Boca Raton, FL, USA :CRC Press/Taylor & Francis Group.
- Davis CB, Markey CE, Busch MA, Busch KW. (2007). Determination of capsaicinoids in habanero peppers by chemometric analysis of UV spectral data. *J. Agric. Food Chem*, 55, 5925–5933.
- Bratt K, Sunnerheim K, Bryngelsson S, Fagerlund A, Engman L, Andersson RE, Dimberg LH. (2003). Avenanthramides in oats (*Avena sativa* L.) and structure-antioxidant activity relationships. *J. Agric. Food Chem*, 51, 594–600.
- Salehi B, Mishra AP, Nigam M, et al. (2018) Resveratrol: A Double-Edged Sword in Health Benefits. *Biomedicines*, 6(3), 91.
- Abe LT, Lajolo FM, Genovese MI. (2012). Potential dietary sources of ellagic acid and other antioxidants among fruits consumed in Brazil: Jaboticaba (*Myrciaria jaboticaba* (Vell.) Berg *Journal of the Science of Food and Agriculture* 92 (8), 1679-1687.
- Rodríguez-García C, Sánchez-Quesada C, Toledo E, Delgado-Rodríguez M, Gaforio JJ. (2019). Naturally Lignan-Rich Foods: A Dietary Tool for Health Promotion? *Molecules*, 24(5), 917.
- Stanić Z. (2017). Curcumin, a Compound from Natural Sources, a True Scientific Challenge - A Review. *Plant Foods Hum Nutr*, 72(1), 1-12.
- Emmanouil H, Papaioannou, Kyriakides ML, Karabelas AJ. (2016). Natural Origin Lycopene and Its “Green” Downstream Processing. *Critical Reviews in Food Science and Nutrition*, 56(4), 686-709.
- Saini R (2011). Coenzyme Q10: The essential nutrient. *J Pharm Bioallied Sci* 3(3), 466–467.

25. Willett WC. (2010). Fruits, vegetables, and cancer prevention: turmoil in the produce section. *J Natl Cancer Inst*, 102, 510–511.
26. Bhattacharyya S, Hossain MSD, Mohanty S, et al. (2010). Curcumin reverses T cell-mediated adaptive immune dysfunctions in tumor-bearing hosts. *Cell Mol Immunol*, 7, 306–315.
27. Lee-Chang C, Bodogai M, Martin-Montalvo A, et al. (2013). Inhibition of breast cancer metastasis by resveratrol-mediated inactivation of tumor-evoked regulatory B cells. *J Immunol*, 191, 4141–4151.
28. Raffoul JJ, Banerjee S, Che M, et al. (2007). Soy isoflavones enhance radiotherapy in a metastatic prostate cancer model. *Int J Cancer*, 120, 2491–2498.
29. Greenlee H., Kwan M.L., Kushi L.H. (2012). Antioxidant supplement use after breast cancer diagnosis and mortality in the LACE cohort. *Cancer*, 118, 2048–2058.
30. Maeda N, Hagihara H, Nakata Y, Hiller S, Wilder J, Reddick R. (2000). Aortic wall damage in mice unable to synthesize ascorbic acid. *Proc Natl Acad Sci U S A*, 97(2), 841–846.
31. Devaraj S, Li D, Jialal I. (1996). The effects of alpha tocopherol supplementation on monocyte function. Decreased lipid oxidation, interleukin 1 beta secretion, and monocyte adhesion to endothelium. *J Clin Invest*, 98(3), 756–763.
32. Romanchik JE, Morel DW, Harrison EH. (1995). Distributions of carotenoids and alpha-tocopherol among lipoproteins do not change when human plasma is incubated in vitro. *J Nutr*, 125(10), 2610–2617.
33. Stephens NG, Parsons A, Schofield PM, Kelly F, Cheeseman K, Mitchinson MJ. (1996). Randomised controlled trial of vitamin E in patients with coronary disease: Cambridge heart antioxidant study (CHAOS). *Lancet*, 347(9004), 781–786.
34. Hozawa A, Jacobs DR Jr, Steffes MW, Gross MD, Steffen LM, Lee DH. (2007). Relationships of circulating carotenoid concentrations with several markers of inflammation, oxidative stress, and endothelial dysfunction: the coronary artery risk development in young adults (CARDIA)/ young adult longitudinal trends in antioxidants (YALTA) study. *Clin. Chem*, 53(3), 447–455.
35. Wannamethee SG, Lowe GD, Rumley A, Bruckdorfer KR, Whincup PH. (2006). Associations of vitamin C status, fruit and vegetable intakes, and markers of inflammation and hemostasis. *Am J Clin. Nutr*, 83(3), 567–574.
36. Quideau S, Deffieux D, Douat-Casassus C, Pouysegu L. (2011). Plant polyphenols: chemical properties, biological activities, and synthesis. *Angew Chem Int Ed Engl*, 50(3), 586–621.
37. Yamakoshi J, Kataoka S, Koga T, Ariga T. (1999). Proanthocyanidin-rich extract from grape seeds attenuates the development of aortic atherosclerosis in cholesterol-fed rabbits. *Atherosclerosis*, 142(1), 139–149.
38. Kaplan M, Hayek T, Raz A, Coleman R, Dornfeld L, Vaya J, et al. (2001). Pomegranate juice supplementation to atherosclerotic mice reduces macrophage lipid peroxidation, cellular cholesterol accumulation and development of atherosclerosis. *J. Nutr*, 131(8), 2082–2089.
39. Joshipura KJ, Hu FB, Manson JE, Stampfer MJ, Rimm EB, Speizer FE, Colditz G, Ascherio A, Rosner B, Spiegelman D, Willett WC. (2001). The effect of fruit and vegetable intake on risk for coronary heart disease. *Ann Int Med*, 134, 1106–1114.
40. Boaz M, Smetana S, Weinstein T, Matas Z, Gafter U, Iaina A, Knecht A, Weissgarten Y, Fainaru M, Green M. (2000). Secondary prevention using antioxidants of cardiovascular disease in endstage renal disease: SPACE. *Eur Heart J*, 21, S458.
41. Kim G. H., Kim J. E., Rhie S. J., Yoon S. (2015). The role of oxidative stress in neurodegenerative diseases. *Exp. Neurobiol*, 24, 325–340.
42. Federico A, Cardaioli E, Da Pozzo P, Formichi P, Gallus GN, Radi E. (2012). Mitochondria, oxidative stress and neurodegeneration. *J Neurol Sci*, 322, 254–262.
43. Masgrau R, Guaza C, Ransohoff RM, Galea E. (2017). “Should we stop saying “glia” and “neuroinflammation”?” *Trends in Molecular Medicine*, 23(6), 486–500.
44. Frank-Cannon TC, Alto LT, McAlpine FE, Tansey MG, (2009). “Does neuroinflammation fan the flame in neurodegenerative diseases?” *Molecular Neurodegeneration*, 4(1), 47.
45. Prakash A, Kumar A. (2014). “Implicating the role of lycopene in restoration of mitochondrial enzymes and BDNF levels in β -amyloid induced Alzheimer’s disease,” *European Journal of Pharmacology*, 741, 104–111.
46. Katayama S, Ogawa H, Nakamura S, (2011). “Apricot carotenoids possess potent anti-amyloidogenic activity in vitro,” *Journal of Agricultural and Food Chemistry*, 59(23), 12691–12696.

47. Suganuma H, Hirano T, Arimoto Y, Inakuma T. (2002). "Effect of tomato intake on striatal monoamine level in a mouse model of experimental Parkinson's disease." *Journal of Nutritional Science and Vitaminology*, 48(3), 251–254
48. Di Matteo V, Pierucci M, Di Giovanni ., et al. (2009). Intake of tomato-enriched diet protects from 6-hydroxydopamine-induced degeneration of rat nigral dopaminergic neurons. *Birth, Life and Death of Dopaminergic Neurons in the Substantia Nigra*, 73, 333–341.
49. Kumar P, Kumar A. (2009). Effect of lycopene and epigallocatechin-3-gallate against 3-nitropropionic acid induced cognitive dysfunction and glutathione depletion in rat: a novel nitric oxide mechanism. *Food and Chemical Toxicology*, 47(10), 2522–2530.
50. Qu M, Li L, Chen C, et al. (2011). Protective effects of lycopene against amyloid β -induced neurotoxicity in cultured rat cortical neurons. *Neuroscience Letters*, 505(3), 286–290.
51. Yi F, Xin H, Wang D. (2013). Lycopene protects against MPP⁺-induced cytotoxicity by maintaining mitochondrial function in SH-SY5Y cells. *Neurochemical Research*, 38(8), 747–757.
52. Xiang S, Liu F, Lin J, et al. (2017). Fucoxanthin inhibits β -amyloid assembly and attenuates β -amyloid oligomer-induced cognitive impairments. *Journal of Agricultural and Food Chemistry*, 65(20), 4092–4102.
53. Lobos P, Bruna B, Cordova A, et al. (2016). Astaxanthin protects primary hippocampal neurons against noxious effects of A β -oligomers. *Neural Plasticity*, 2016, 3456783
54. Tiribuzi R, Crispolti L, Chiurchiù V, et al. (2017). Trans-Crocetin improves amyloid- β degradation in monocytes from Alzheimer's disease patients. *Journal of the Neurological Sciences*, 372, 408–412.
55. Ye Q, Zhang X, Huang B, Zhu Y, Chen X. (2013). Astaxanthin suppresses MPP⁺-induced oxidative damage in PC12 cells through a Sp1/NR1 signaling pathway. *Marine Drugs*, 11(12), 1019–1034.
56. Isonaka R, Hiruma H, Katakura T, Kawakami T. (2011). Inhibition of superoxide dismutase selectively suppresses growth of rat spinal motor neurons: comparison with phosphorylated neurofilament-containing spinal neurons. *Brain Research*, 1425, 13–19
57. Balmus I, Ciobica A, Trifan A, Stanciu C. (2016). The implications of oxidative stress and antioxidant therapies in inflammatory bowel disease: clinical aspects and animal models. *Saudi Journal of Gastroenterology*, 22(1), 3–17.
58. Mete R, Tulubas F, Oran M, Yılmaz A, Avcı BA, Yildiz K, Turan CB, Gurel A. (2013). The role of oxidants and reactive nitrogen species in irritable bowel syndrome: A potential etiological explanation. *Med Sci Monit*, 19, 762–766.
59. Lang A, Salomon N, Wu JC, et al. (2015). Curcumin in combination with mesalamine induces remission in patients with mild-to-moderate ulcerative colitis in a randomized controlled trial. *Clin Gastroenterol Hepatol*, 13, 1444–1449
60. Gerhardt H, Seifert F, Buvari P, et al. (2001). Therapy of active Crohn disease with *Boswellia serrata* extract H 15 (in German) *Z Gastroenterol*, 39, 11–17.
61. Krebs S, Omer TN, Omer B. (2010). Wormwood (*Artemisia absinthium*) suppresses tumour necrosis factor alpha and accelerates healing in patients with Crohn's disease - a controlled clinical trial. *Phytomedicine*, 17, 305–309
62. Zhu W, Li Y, Gong J, et al. (2015). *Tripterygium wilfordii* Hook. f. versus azathioprine for prevention of postoperative recurrence in patients with Crohn's disease: a randomized clinical trial. *Dig Liver Dis*, 47, 14–19
63. Feagan BG, Sandborn WJ, Mittmann U, et al. (2008). Omega-3 free fatty acids for the maintenance of remission in Crohn disease: the EPIC Randomized Controlled Trials. *JAMA*, 299, 1690–1697
64. Langhorst J, Frede A, Knott M, et al. (2014). Distinct kinetics in the frequency of peripheral CD4⁺ T cells in patients with ulcerative colitis experiencing a flare during treatment with mesalazine or with a herbal preparation of myrrh, chamomile, and coffee charcoal. *PLoS One*, 9, e104257
65. Barbosa DS, Cecchini R, El Kadri MZ, et al. (2003). Decreased oxidative stress in patients with ulcerative colitis supplemented with fish oil omega-3 fatty acids. *Nutrition*, 19, 837–842
66. Jadeja RN, Devkar RV, Srinivas N. (2017). Oxidative stress in liver diseases: pathogenesis, prevention, and therapeutics. *Oxid Med Cell Longev*, 2017, 1–2
67. Wang ME, Chen YC, Chen IS, Hsieh SC, Chen SS, Chiu CH. (2012). Curcumin protects against thioacetamide-induced hepatic fibrosis by attenuating the inflammatory response and inducing apoptosis of damaged hepatocytes. *J Nutr Biochem*, 23, 1352–1366

68. Reyes-Gordillo K, Segovia J, Shibayama M, Vergara P, Moreno MG, Muriel P. (2007). Curcumin protects against acute liver damage in the rat by inhibiting NF-kappaB, proinflammatory cytokines production and oxidative stress. *Biochim Biophys Acta*, 1770, 989–996
69. Hernández-Ortega LD, Alcántar-Díaz BE, Ruiz-Corro LA, Sandoval-Rodríguez A, Bueno-Topete M, Armendariz-Borunda J, Salazar-Montes AM. (2012). Quercetin improves hepatic fibrosis reducing hepatic stellate cells and regulating pro-fibrogenic/anti-fibrogenic molecules balance. *J Gastroenterol Hepatol*, 27, 1865–1872.
70. Kasdallah-Grissa A, Mornagui B, Aouani E, Hammami M, Gharbi N, Kamoun A, El-Fazaa S. (2006). Protective effect of resveratrol on ethanol-induced lipid peroxidation in rats. *Alcohol Alcohol*, 41, 236–239.
71. Grievink L, Smit HA, Ocké MC, Van't VP, Kromhout D. (1998). Dietary intake of antioxidant (pro)-vitamins, respiratory symptoms and pulmonary function: the MORGEN study. *Thorax*, 53, 166–171.
72. Santus P, Sola A, Carlucci P. (2005). Lipid peroxidation and 5-lipoxygenase activity in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*, 171, 838–843.
73. Rautalahti M, Virtamo J, Haukka J, Heinonen OP, Sundvall J, et al. (1997). The effect of alpha-tocopherol and beta-carotene supplementation on COPD symptoms. *Am J Respir Crit Care Med*, 156, 1447–1452.
74. Sargeant LA, Jaeckel A, Wareham NJ. (2000). Interaction of vitamin C with the relation between smoking and obstructive airway disease in EPOC Norfolk. European Prospective Investigation into Cancer and Nutrition. *Eur Respir J*, 16, 397–403.
75. Kode A, Rajendrasozhan S, Caito S, Yang SR, Megson IL. (2008). Resveratrol induces glutathione synthesis by activation of Nrf2 and protects against cigarette smoke mediated oxidative stress in human lung epithelial cells. *Am J Physiol Lung Cell Mol Physiol*, 294, L478–L488.
76. Biswas SK, McClure D, Jimenez LA, Megson IL, Rahaman I. (2005). Curcumin induces glutathione biosynthesis and inhibits NF-kappaB activation and interleukin-8 release in alveolar epithelial cells: mechanism of free radicals scavenging activity. *Antioxid Redox Signal*, 7, 32–41.
77. Hong JY, Lee CY, Lee MG, Kim YS. (2018). Effects of dietary antioxidant vitamins on lung functions according to gender and smoking status in Korea: a population-based cross-sectional study. *BMJ Open*, 8(4), e020656.
78. Fitzpatrick AM, Teague WG, Holguin F, Yeh M, Brown LA. (2009). Airway glutathione homeostasis is altered in children with severe asthma: evidence for oxidant stress. *J Allergy Clin Immunol*, 12, 146–152.
79. Nurmatov U, Devereux G, Sheikh A. (2011). Nutrients and foods for the primary prevention of asthma and allergy: systematic review and meta-analysis. *J Allergy Clin Immunol*, 127, 724–733.
80. Pearson PJ, Lewis SA, Britton J, Fogarty A. (2004). Vitamin E supplements in asthma: a parallel group randomised placebo controlled trial. *Thorax*, 59, 652–656.
81. Kurti SP, Murphy JD, Ferguson CS, Brown KR, Smith JR, Harms CA. (2016). Improved lung function following dietary antioxidant supplementation in exercise-induced asthmatics. *Respir Physiol Neurobiol*, 220, 95–101.
82. Lovett-Racke AE, Racke MK. (2002). Retinoic acid promotes the development of Th2-like human myelin basic protein-reactive T cells. *Cell Immunol*, 215, 54–60.
83. Romieu I, Varraso R, Avenel V, Leynaert B, Kauffmann F, Clavel-Chapelon F. (2006). Fruit and vegetable intakes and asthma in the E3N study. *Thorax*, 61, 209–215.